

Terrestrial analogues for extraterrestrial life

I. N. Reid

*Maryland Astrobiology Consortium
Space Telescope Science Institute
3700 San Martin Drive
USA
inr@stsci.edu*

; K.R. Sowers, H.D. Shukla, S. MacAuley, T. Miller, R. Suvanasuthi, R. Belas, A. Colman, F.T. Robb,
P. DasSarma, J.A. Müller, F. Chen, R. Cavicchioli, S. DasSarma

*Maryland Astrobiology Consortium
Center of Marine Biotechnology, University of Maryland)
USA*

W.B. Sparks, S. Lubow, M. McGrath, M. Livio, J. Valenti

*Maryland Astrobiology Consortium
Space Telescope Science Institute
3700 San Martin Drive
USA*

The past decade has seen the discovery of planetary companions to ~120 stellar neighbours. Those planets are subject to a wide range of environments, from hot Jupiters at ambient temperatures >1400K to gas giants in Jupiter-like orbits at $T_{\text{amb}} \sim 150$ K. A key issue facing astrobiology is assessing what subset of those environments might be hospitable to life; in particular, since cold environments are much more common than hot, what are the low-temperature limits for species growth and survival?

We have been conducting experiments designed to test the cold limits for growth of four extremophile species: two methanogens, *Methanosarcina acetivorans*, a mesophile, and *Methanococcoides burtonii*, an Antarctic psychrotroph; and two extremely halophilic archaea, *Halobacterium* sp. NRC-1, a mesophile, and *Halorubrum lacusprofundi*, an Antarctic psychrotroph. Both types of species are potentially relevant to life in Martian environments: methanogens are one of the proposed sources of significant methane in the present-day Martian atmosphere, while halophiles could take advantage of sub-surface brine, which remains liquid at temperatures well below 270K. Initial results show growth of the psychrotrophic archaea at sub-zero temperatures. Both types of organisms feature an apparent morphological response to cold environments.